

Order-Types, Trees, and a  
Problem of Erdős and Hajnal

by

Keith J. Devlin

(Oslo)

§0. Introduction

We work in ZFC set theory throughout, and use the usual notation and conventions. In particular, an ordinal is the set of all its predecessors and a cardinal is an initial ordinal. The cardinality of a set  $X$  is denoted by  $|X|$ . As usual, if  $\sigma, \rho$  are order types,  $\sigma \leq \rho$  denotes that there is an order preserving monomorphism  $F: \sigma \rightarrow \rho$ . (We use  $\sigma, \rho$  to denote order types, all other lower case Greek letters denoting ordinals, with  $\kappa, \lambda$ , in particular, for cardinals). We assume some familiarity with the notion of constructibility, as in, say, [1]. In particular, we assume the reader knows the usual proof that  $V = L \rightarrow \text{GCH}$  (see [1].) We use  $L_\alpha$ ,  $\alpha \in \text{On}$ , to denote the levels in the constructible hierarchy.

Let  $\Phi$  denote the following proposition: Whenever  $\rho$  is an order type of cardinality  $\omega_2$  such that  $\omega_2 \leq \rho$  and  $\omega_2^* \not\leq \rho$  there is  $\sigma \subset \rho$  of cardinality  $\omega_1$  such that  $\omega_1 \leq \sigma$  and  $\omega_1^* \not\leq \sigma$ . Without assuming GCH, the status of  $\Phi$  is of little interest, of course, since a set of reals of cardinality  $\omega_2$  embeds neither  $\omega_2$ ,  $\omega_2^*$  nor  $\omega_1$ ,  $\omega_1^*$ . In [2] (§ 7, Problem I), Erdős and Hajnal ask what happens to  $\Phi$  if we do assume GCH. In this paper, we prove that if  $V = L$ , then  $\neg \Phi$ . The proof uses Jensen's answer to an earlier question of ours concerning subtrees of  $\omega_1$ -trees, and we end the paper with a brief discussion of this topic, and a remark concerning the consistency of  $\Phi$ .

§1.  $V=L \rightarrow \exists T$

A tree is a poset  $\mathbb{T} = \langle T, \leq_T \rangle$  such that for any  $x \in T$ ,  $\{y \in T \mid y <_T x\}$  is well ordered by  $\leq_T$ . (For such  $\mathbb{T}$ ,  $x$ , we call the order type of this set (under  $<_T$ ) the height of  $x$  in  $\mathbb{T}$ .) For any ordinal  $\alpha$ , we set  $T_\alpha = \{x \in T \mid x \text{ has height } \alpha \text{ in } \mathbb{T}\}$ , the  $\alpha$ 'th level of  $\mathbb{T}$ . We write  $\mathbb{T} \restriction \alpha$  for  $\langle \bigcup_{\beta < \alpha} T_\beta, \leq_T \restriction (\bigcup_{\beta < \alpha} T_\beta) \rangle$ . A branch of  $\mathbb{T}$  is a maximal linearly ordered subset of  $T$ . If it has order type  $\alpha$  it is an  $\alpha$ -branch.

Let  $\theta$  be an ordinal,  $\kappa$  a cardinal.  $\mathbb{T}$  is a  $(\theta, \kappa)$ -tree iff  $(\forall \alpha < \theta)(T_\alpha \neq \emptyset \ \& \ |T_\alpha| < \kappa) \ \& \ T_\theta = \emptyset$ .  $\mathbb{T}$  is normal iff whenever  $\alpha < \beta < \theta$  and  $x \in T_\alpha$  there are distinct  $y, y' \in T_\beta$  with  $x <_T y, y'$ . A tree  $\mathbb{T}$  is  $\kappa$ -Aronszajn iff it is a  $(\kappa, \kappa)$ -tree with no  $\kappa$ -branches.  $\mathbb{T}$  is  $\kappa$ -Kurepa iff it is a  $(\kappa, \kappa)$ -tree with at least  $\kappa^+$   $\kappa$ -branches. In both cases, we do not bother to mention the  $\kappa$  if  $\kappa = \omega_1$ . (We ignore  $\kappa = \omega$  from the outset.)

Solovay has proved that if  $V = L$ , there is a Kurepa tree. For our purposes, a somewhat stronger result is required.

Let us begin by proving three fundamental lemmas on constructibility.

Lemma 1

Assume  $V = L$ . If  $M \prec L_{\omega_1}$ , then  $M = L_\alpha$  for some  $\alpha \leq \omega_1$ .

Proof: By absoluteness considerations, it suffices to show that  $M$  is transitive. Let  $x \in L_{\omega_1}$ . Then  $x$  is countable. Now,  $L_{\omega_1}$  has an  $L_{\omega_1}$ -definable well ordering,  $<_L$ . Let  $J_x$  be the  $<_L$ -least bijection  $J_x: \omega \leftrightarrow x$ . Then  $J_x \in L_{\omega_1}$  and is  $L_{\omega_1}$ -definable from  $x$ . Hence,  $x \in M \rightarrow J_x \in M$ . But clearly,  $\omega \subset M$ . Thus  $x \in M \rightarrow J_x''\omega \subset M \rightarrow x \subset M$ , as required.

### Lemma 2

Assume  $V = L$ . If  $M \prec L_{\omega_2}$ , then  $M \cap L_{\omega_1} = L_\alpha$  for some  $\alpha \leq \omega_1$ .

Proof:  $L_{\omega_1}$  is  $L_{\omega_2}$ -definable, so clearly  $M \cap L_{\omega_1} \prec L_{\omega_1}$ . By lemma 1,  $M \cap L_{\omega_1} = L_\alpha$  for some  $\alpha \leq \omega_1$ .

### Lemma 3

Let  $\beta > \alpha > \omega$ ,  $p \in L_\beta$ . Suppose that  $L_{\beta+1} \models "\alpha \text{ is a regular uncountable cardinal}"$  and that  $L_\beta$  is the smallest  $X \prec L_\beta$  such that  $p \in X$  and  $X \cap \alpha$  is transitive. Then there is an  $L_{\beta+1}$ -definable map of  $\omega$  cofinally into  $\alpha$ .

Proof: Let us use the notation  $X \prec_{\Sigma_n} L_\beta$  to mean that  $X$  is an elementary substructure of  $\langle L_\beta, \epsilon \rangle$  when we restrict our attention to the  $\Sigma_n$  formulas of set theory only. For each  $n \in \omega$ , let  $X_n$  be the smallest  $X \prec_{\Sigma_n} L_\beta$  such that  $p \in X$  and  $X \cap \alpha$  is transitive. Thus  $X_n$  is  $L_\beta$ -definable. But  $L_{\beta+1} \models "\alpha \text{ is a regular uncountable cardinal}"$ , so clearly  $X_n \cap \alpha \in \alpha$ . Let  $\alpha_n = X_n \cap \alpha$ . Since  $\langle X_n \mid n < \omega \rangle$  is  $L_{\beta+1}$ -definable, so is  $\langle \alpha_n \mid n < \omega \rangle$ . But  $\bigcup_{n < \omega} X_n = L_\beta$  by assumption on  $L_\beta$ , so  $\sup_{n < \omega} \alpha_n = \alpha$ , and we are done.

The following theorem was proved by Jensen in answer to an old question of ours.

### Theorem 4 (Jensen)

Assume  $V = L$ . Then there is a Kurepa tree no subset of which is (under the inherited ordering) an Aronszajn tree.

Proof: Define a function  $h: \omega_1 \rightarrow \omega_1$  by setting  $h(\alpha) =$  the least  $\gamma$  such that  $L_{\gamma+1} \models "\alpha \text{ is countable}"$ . We define an  $(\omega_1, \omega_1)$ -tree  $\mathbb{T}$  by induction on the levels. The elements of  $\mathbb{T}$  will be countable ordinals, and we have  $\alpha <_{\mathbb{T}} \beta \rightarrow \alpha < \beta$ .

Each  $\mathbb{T} \restriction \alpha$  will be a normal  $(\alpha, \omega_1)$ -tree.

As we proceed, we simultaneously define a function  $f: \omega_1 \rightarrow \omega_1$  by setting  $f(\alpha) =$  the least  $\gamma$  such that  $\alpha, \mathbb{T} \restriction \alpha \in L_\gamma \setminus L_{\omega_1}$  (by lemma 1).

Set  $T_0 = \{0\}$ . If  $T_\alpha$  is defined,  $T_{\alpha+1}$  is the result of appointing (in a minimal way) two new ordinals to succeed each member of  $T_\alpha$ . We are left with the definition of  $T_\alpha$  when  $\lim(\alpha)$  and  $\mathbb{T} \restriction \alpha$  is defined, and is a normal  $(\alpha, \omega_1)$ -tree.

Let  $S(\alpha)$  be the set of all  $U \subset \mathbb{T} \restriction \alpha$  such that:

- (i)  $U \in \bigcup_{\xi < h(\alpha)} L_\xi$ ;
- (ii)  $U$  is a normal  $(\alpha, \omega_1)$ -tree (under the inherited ordering);
- (iii)  $U$  is thin in  $\mathbb{T} \restriction \alpha$  (i.e. for any  $x \in U$  there is a  $y \in \mathbb{T} \restriction \alpha - U$  such that  $x <_{\mathbb{T}} y$ );
- (iv)  $L_{h(\alpha)} \models "U \text{ is an Aronszajn tree}"$ .

We let  $T_\alpha$  consist of (minimally appointed ordinals as) one point extensions of each  $\alpha$ -branch  $b$  of  $\mathbb{T} \restriction \alpha$  such that  $b \in L_{f(\alpha)}$  and  $b \not\subset U$  for any  $U \in S(\alpha)$ . By condition (iii) above, it is clear that (since  $L_{f(\alpha)} \models "\alpha$  and  $S(\alpha)$  are countable")  $\mathbb{T} \restriction \alpha + 1$  is a normal  $(\alpha + 1, \omega_1)$ -tree.

Set  $\mathbb{T} = \bigcup_{\alpha < \omega_1} \mathbb{T} \restriction \alpha$ , a normal  $(\omega_1, \omega_1)$ -tree.

We prove that  $\mathbb{T}$  is Kurepa. Suppose not. Then we may let  $\langle c_\alpha \mid \alpha < \omega_1 \rangle$  be the  $<_L$ -least enumeration of all the  $\omega_1$ -branches of  $\mathbb{T}$ .  $\mathbb{T}$  is clearly  $L_{\omega_2}$ -definable, and hence so is  $\langle c_\alpha \mid \alpha < \omega_1 \rangle$ . Using this fact, we define an  $\omega_1$ -branch  $b$  of  $\mathbb{T}$  such that  $b \neq c_\alpha$  for all  $\alpha < \omega_1$ , giving the required result.

Define a chain of submodels  $X_0 \prec X_1 \prec \dots \prec X_\nu \prec \dots \prec L_{\omega_2}$  ( $\nu < \omega_1$ ) as follows:

$X_0 = \text{the smallest } X \in L_{\omega_2};$

$X_{\nu+1} = \text{the smallest } X \in L_{\omega_2} \text{ such that } X_\nu \cup \{X_\nu\} \subset X;$

$X_\gamma = \bigcup_{\nu < \gamma} X_\nu \text{ if } \lim(\gamma).$

For each  $\nu < \omega_1$ , let  $\pi_\nu: X_\nu \cong L_{\psi(\nu)}$ . By lemma 2,

$\pi_\nu(\omega_1) = \omega_1 \cap X_\nu = \alpha_\nu$ , say. (Note that  $\langle \alpha_\nu | \nu < \omega_1 \rangle$  is a continuous sequence). Hence  $\pi_\nu(\mathbb{T}) = \mathbb{T} \restriction \alpha_\nu$  and  $\pi_\nu(\langle c_\alpha | \alpha < \omega_1 \rangle) = \langle c_\alpha \cap \alpha_\nu | \alpha_\nu \rangle$ .

Also,  $L_f(\alpha_\nu) \models "\alpha_\nu \text{ is countable}"$  whereas  $L_{\psi(\nu)} \models "\alpha_\nu = \omega_1"$ , so  $\psi(\nu) < f(\alpha_\nu)$  for each  $\nu < \omega_1$ . It follows that  $\langle c_\alpha \cap \alpha_\nu | \alpha < \alpha_\nu \rangle \in L_f(\alpha_\nu)$ .

We define, by induction, a sequence  $b_0 \subset \dots \subset b_\nu \subset \dots (\nu < \omega_1)$  such that, for each  $\nu < \omega_1$ ,  $b_\nu$  is an  $\alpha_\nu$ -branch of  $\mathbb{T} \restriction \alpha_\nu$ , with an extension on level  $\alpha_\nu$  of  $\mathbb{T}$ , such that  $\alpha < \alpha_\nu \rightarrow b_\nu \neq c_\alpha \cap \alpha_\nu$ . We shall then just set  $b = \bigcup_{\nu < \omega_1} b_\nu$  and be done, of course.

To define  $b_0$ , observe that  $\langle c_\alpha \cap \alpha_0 | \alpha < \alpha_0 \rangle, S(\alpha_0), \alpha_0$ ,  $\mathbb{T} \restriction \alpha_0 \in L_f(\alpha_0)$  and  $L_f(\alpha_0) \models "\alpha_0, S(\alpha_0), \mathbb{T} \restriction \alpha_0 \text{ are countable}"$ . Hence, working inside  $L_f(\alpha_0)$ , we may let  $b_0$  be the  $<_L$ -least  $\alpha_0$ -branch of  $\mathbb{T} \restriction \alpha_0$  such that  $b_0 \not\subset U$  for any  $U \in S(\alpha_0)$  and  $\alpha < \alpha_0 \rightarrow b_0 \neq c_\alpha \cap \alpha_0$ . Then, since  $S(\alpha_0) \subset L_f(\alpha_0)$ ,  $b_0$  has an extension on  $T_{\alpha_0}$ .

If  $b_0 \subset \dots \subset b_\nu$  are already (suitably) defined,  $b_{\nu+1}$  is defined similarly (to contain the  $T_{\alpha_\nu}$ -extension of  $b_\nu$ , of course).

Finally, suppose  $\lim(\nu)$  and that  $b_0 \subset b_1 \subset \dots \subset b_\xi \subset \dots (\xi < \nu)$  are suitably defined. Let  $b_\nu = \bigcup_{\xi < \nu} b_\xi$ . Since  $\langle \alpha_\nu | \nu < \omega_1 \rangle$  is continuous,  $b_\nu$  is an  $\alpha_\nu$ -branch of  $\mathbb{T} \restriction \alpha_\nu$  and  $\alpha < \alpha_\nu \rightarrow b_\nu \neq c_\alpha \cap \alpha_\nu$ . We must show that  $b_\nu$  extends on  $T_{\alpha_\nu}$ . We show first that  $b_\nu \in L_f(\alpha_\nu)$ . Clearly,  $b_\nu$  is definable from  $\mathbb{T} \restriction \alpha_\nu, \langle \alpha_\xi | \xi < \nu \rangle, \langle S(\alpha_\xi) | \xi < \nu \rangle, \langle c_\alpha \cap \alpha_\nu | \alpha < \alpha_\nu \rangle$ , so it reduces to proving that  $\langle \alpha_\xi | \xi < \nu \rangle \in L_f(\alpha_\nu)$ .

And since  $\alpha_\xi = \omega_1^{L_{\psi(\xi)}}$  for each  $\xi$ , this will be so providing

$\langle L_{\psi}(\xi) \mid \xi < \nu \rangle \in L_{f(\alpha_\nu)}$ . Now,  $\psi(\nu) < f(\alpha_\nu)$ , so  $L_{\psi(\nu)} \in L_{f(\alpha_\nu)}$ . Thus, working inside  $L_{f(\alpha_\nu)}$ , we may define a chain  $Y_0 \prec Y_1 \prec \dots \prec Y_\xi \prec \dots \prec L_{\psi(\nu)}$  ( $\xi < \nu$ ) exactly as  $\langle X_\xi \mid \xi < \omega_1 \rangle$  was defined from  $L_{\omega_2}$ . But look, in defining  $\langle X_\xi \mid \xi < \nu \rangle$ , we could equally well have used  $X_\nu$  in place of  $L_{\omega_2}$  (since  $\xi < \nu \rightarrow X_\xi \prec X_\nu \prec L_{\omega_2}$ ). Then, since  $X_\nu \cong L_{\psi(\nu)}$ , an easy induction argument shows that  $X_\xi \cong Y_\xi$  for all  $\xi < \nu$ . Hence  $Y_\xi \cong L_{\psi}(\xi)$ ,  $\xi < \nu$ . But  $\langle Y_\xi \mid \xi < \nu \rangle \in L_{f(\alpha_\nu)}$ , so  $\langle L_{\psi}(\xi) \mid \xi < \nu \rangle \in L_{f(\alpha_\nu)}$ , as required.

Thus,  $b_\nu$  will be proved to extend on level  $\alpha_\nu$  if we can show that for any  $U \in S(\alpha_\nu)$ ,  $b_\nu \not\subseteq U$ . To this end, note first that our above argument did more than prove that

$b_\nu \in L_{f(\alpha_\nu)}$ . What we actually proved was that  $b_\nu$  is  $L_{\psi(\nu)+1}$ -definable (from elements of  $L_{\psi(\nu)+1}$ .) Now,  $\alpha_\nu$  is uncountable in  $L_{\psi(\nu)+1}$  but countable in  $L_{h(\alpha_\nu)+1}$ . (For the former, note that  $\pi_\nu^{-1} : L_{\psi(\nu)} \prec L_{\omega_2}$  and  $\pi_\nu^{-1}(\alpha_\nu) = \omega_1$ .) Hence  $\psi(\nu) < h(\alpha_\nu)$ .

Case A:  $\psi(\nu) + 1 < h(\alpha_\nu)$ .

Then  $b_\nu \in L_{h(\alpha_\nu)}$ . So as  $L_{h(\alpha_\nu)} \models "U \text{ is Aronszajn}"$  for any  $U \in S(\alpha_\nu)$ , we cannot have  $b_\nu \subseteq U$  for any such  $U$ .

Case B:  $\psi(\nu) + 1 = h(\alpha_\nu)$ .

Thus  $S(\alpha_\nu) \subseteq L_{\psi(\nu)}$ . Let  $U \in S(\alpha_\nu)$ . For  $i < \tau \leq \nu$ , set  $\pi_{i\tau} = \pi_\tau \circ \pi_i^{-1}$ . Then  $\langle L_{\psi(\nu)}, (\pi_{\tau\nu})_{\tau < \nu} \rangle$  is (isomorphic to) the direct limit of the elementary system  $\langle (L_{\psi(\tau)})_{\tau < \nu}, (\pi_{i\tau})_{i < \tau < \nu} \rangle$ . So for some  $\tau < \nu$  and some  $U' \in L_{\psi(\tau)}$ ,  $U = \pi_{\tau\nu}(U')$ . Let  $\tau$  be the least such. It is then easily observed that  $L_{\psi(\tau)}$  is the smallest  $X \prec L_{\psi(\tau)}$  such that  $U' \in L_{\psi(\tau)}$  and  $X \cap \alpha_\tau$  is transitive. Also,  $L_{\psi(\tau)} + 1 \models " \alpha_\tau \text{ is a regular uncountable cardinal} "$ . (By applying  $\pi_\tau^{-1}$ ).

Thus by lemma 3, it follows that  $\alpha_\tau$  is countable in  $L_{\psi(\tau)+2}$ . Hence  $\psi(\tau) + 1 = h(\alpha_\tau)$ . But look, there is no  $L_{\psi(\nu)}$ -definable  $\alpha_\nu$ -branch of  $U$ . (By applying  $\pi_\nu^{-1}$ ). Hence there can be no  $L_{\psi(\tau)}$ -definable  $\alpha_\tau$ -branch of  $U'$ . Thus  $L_{h(\alpha_\tau)} \models "U' \text{ is Aronszajn}"$ . It follows readily that  $U' \in S(\alpha_\tau)$ . Hence  $b_\tau \notin U'$ . But  $U' = U \cap \alpha_\tau$  and  $b_\tau = b_\nu \cap \alpha_\tau$ . Hence  $b_\nu \notin U$ .

This completes the proof that  $\mathbb{T}$  is Kurepa.

Now suppose that  $\mathbb{T}$  has an Aronszajn subtree,  $U$ . It is easily seen that there is no loss of generality in assuming that  $U$  is normal and thin in  $\mathbb{T}$ . We may also assume that  $U$  is the  $<_L$ -least such. Hence  $U$  is  $L_{\omega_2}$ -definable. So  $U \in X_0$ . Clearly,  $U' = \pi_0(U) = U \cap \mathbb{T} \restriction \alpha_0$ . As  $U$  is Aronszajn, there is no  $L_{\omega_2}$ -definable  $\omega_1$ -branch of  $U$ . Thus  $L_{\psi(0)+1} \models "U' \text{ is Aronszajn}"$ . But, using lemma 3, it is immediately clear that  $\psi(0) + 1 = h(\alpha_0)$ . Hence we see that  $U' \in S(\alpha_0)$ . But by construction, no  $\alpha_0$ -branch of  $U'$  ever extended on  $T_{\alpha_0}$ , so how ever can  $U \supset U'$  be cofinal in  $\omega_1$ ? This contradiction completes the proof.  $\blacksquare$

The above theorem, together with our next result, shows that  $V = L \rightarrow \neg \Phi$ .

### Theorem 5

Assume GCH. Suppose that there is a Kurepa tree  $\mathbb{T}$ , no subset of which is an Aronszajn tree. Then  $\neg \Phi$ .

Proof: Without loss of generality, we may assume that  $T \subset \cup_{\alpha < \omega_1} 2^\alpha$  and that  $s \leq_T t$  iff  $s$  is an initial segment of  $t$  (written  $s \text{ inl } t$ ).

Let  $\rho$  be the set of all  $\omega_1$ -branches of  $\mathbb{T}$ , ordered lexicographically. Thus  $|\rho| = \omega_2$ . We show that  $\omega_2, \omega_2^* \not\leq \rho$ .

Let  $D = \{s \in \bigcup_{\alpha < \omega_1} 2^\alpha \mid (\exists b \in \rho)(s \text{ inl } b)\}$ . Then  $|D| = \omega_1$  by GCH.

Suppose  $\omega_2 \leq \rho$ , and let  $\langle b_\nu \mid \nu < \omega_2 \rangle$  be an  $\omega_2$ -sequence from  $\rho$ . Let  $A = \{\nu \in \omega_2 \mid \lim(\nu)\}$ . By induction on  $\nu \in A$ , pick  $s_\nu \in D$  such that  $\tau \in A \cap \nu \rightarrow s_\tau \neq s_\nu$  (By demanding that  $s_\nu \text{ inl } b_\nu$  but  $\neg s_\nu \text{ inl } b_{\nu+1}$  for each  $\nu \in A$ ). Then  $\{s_\nu \mid \nu \in A\}$  is a set of  $\omega_2$  distinct members of  $D$ , which is absurd. Similarly if  $\omega_2^* \leq \rho$ .

Now let  $\sigma \subset \rho$ ,  $|\sigma| = \omega_1$ . We show that  $\omega_1 \leq \sigma$  or  $\omega_1^* \leq \sigma$ . Suppose  $\rho = \langle \rho, \rightarrow \rangle$ .

Let  $U = \{s \in \bigcup_{\alpha < \omega_1} 2^\alpha \mid (\exists b \in \sigma)(s \text{ inl } b)\}$ . Thus  $U \subset \mathbb{T}$ .

Case 1: For some  $b \in \sigma$  it is the case that whenever  $s \text{ inl } b$  there is  $b' \in \sigma$  such that  $s \text{ inl } b'$  and  $b' \neq b$ . By induction, pick a strictly  $<_{\mathbb{T}}$ -increasing sequence  $\langle s_\nu \mid \nu < \omega_1 \rangle$  of initial sections of  $b$ , and a pairwise distinct sequence  $\langle b_\nu \mid \nu < \omega_1 \rangle$  of members of  $\sigma - \{b\}$  such that  $s_\nu \text{ inl } b_\nu$ .

Then either  $\{b_\nu \mid \nu < \omega_1 \ \& \ b_\nu \rightarrow b\}$  or else  $\{b_\nu \mid \nu < \omega_1 \ \& \ b \rightarrow b_\nu\}$  has cardinality  $\omega_1$ . But  $\langle b_\nu \mid \nu < \omega_1 \ \& \ b_\nu \rightarrow b \rangle$  is a  $\rightarrow$ -increasing sequence from  $\sigma$  and  $\langle b_\nu \mid \nu < \omega_1 \ \& \ b \rightarrow b_\nu \rangle$  is a  $\rightarrow$ -decreasing sequence from  $\sigma$ .

Case 2: Otherwise.

Then, for each  $b \in \sigma$  there is  $s_b \in U$  such that  $s_b \text{ inl } b$  and for all  $b' \in \sigma$ ,  $s_b \text{ inl } b' \rightarrow b' = b$ . Let  $U' = \{s \mid (\exists b \in \sigma)(s \text{ inl } s_b)\}$ . Then  $U' \subset U$ . We know that  $U'$  cannot be an Aronszajn tree. And yet  $U'$  is an  $(\omega_1, \omega_1)$ -tree. Hence  $U'$  has an  $\omega_1$ -branch,  $d$ . For  $b_1, b_2 \in \sigma$ ,  $s_{b_1}$  and  $s_{b_2}$  must be  $<_{\mathbb{T}}$ -incomparable. Hence for each  $s \text{ inl } d$  there must be a  $b \in \sigma$  such that  $s \text{ inl } s_b$ .



and  $\neg s_b \text{ inl } d$ . So, for each  $s \text{ inl } d$  there is  $b \in \sigma$  such that  $s \text{ inl } b$  and  $b \neq d$ . So, as in Case 1,  $\omega_1 \leq \sigma$  or  $\omega_1^* \leq \sigma$ .  $\square$

### §3. Subtrees of $\omega_1$ -trees.

We saw above that it is consistent that there is a Kurepa tree with no Aronszajn subtree. Since the existence of Kurepa trees is not provable in ZFC (see [4]), we could not hope to eliminate the use of  $V = L$  in establishing that result. However, in ZFC, it is possible to construct a normal  $(\omega_1, \omega_1)$ -tree with no Aronszajn subtree. In fact, we have:

#### Theorem 6.

there is a normal  $(\omega_1, \omega_1)$ -tree  $\mathbb{T}$  such that:

- (i)  $\mathbb{T}$  has no Aronszajn subtree.
- (ii) if  $\mathbb{T}'$  is any normal  $(\omega_1, \omega_1)$ -tree then either  $\mathbb{T}'$  has an Aronszajn subtree or else  $\mathbb{T} \leq \mathbb{T}'$ .

Proof: Let  $T = \{s \in 2^{\omega_1} \mid |\{\alpha \in \omega_1 \mid s(\alpha) = 1\}| < \omega\}$ , and make  $T$  into a tree by setting  $s \leq t$  iff  $s \subseteq t$ . Clearly,  $\mathbb{T}$  is a normal  $(\omega_1, \omega_1)$ -tree such that every point in  $T$  lies on an  $\omega_1$ -branch of  $\mathbb{T}$ .

- (i) Suppose  $U \subseteq T$ ,  $U$  a normal  $(\omega_1, \omega_1)$ -tree. We show that  $U$  is not an Aronszajn tree. It is easily seen that we may assume that  $U$  is an initial segment of  $\mathbb{T}$ . (If the initialisation of  $U$  in  $\mathbb{T}$  has an  $\omega_1$ -branch, so must  $U$  itself!) Set  $C = \{\alpha \in \omega_1 \mid \lim(\alpha)\}$ . For each  $\alpha \in C$ , let  $s_\alpha \in U$  be arbitrary. For each  $\alpha \in C$ , define  $f(\alpha) =$  the largest  $\beta < \alpha$  such that  $s_\alpha(\beta) = 1$ , or else  $f(\alpha) = 0$  if no such  $\beta$  exists. Then  $f: C \rightarrow \omega_1$  is regressive, so by a well known

theorem of Fodor (see [1], Chapter 3, for example) we can find stationary set  $X \subseteq C$  such that  $f''X = \{\beta_0\}$  for some fixed  $\beta_0$ . It follows immediately that there must be an uncountable set  $Y \subseteq X$  such that  $\alpha, \beta \in Y \ \& \ \alpha < \beta \rightarrow s_\alpha \leq_T s_\beta$ . Hence  $\{s_\alpha \mid \alpha \in Y\}$  determines an  $\omega_1$ -branch of  $U$ .

(ii) Let  $\mathcal{T}$  be a normal  $(\omega_1, \omega_1)$ -tree with no Aronszajn subtree.

By replaing  $\mathcal{T}'$  by a subtree if necessary, we assume  $\mathcal{T}'$  is such that every point in  $\mathcal{T}'$  has exactly two distinct immediate successors. But look, as  $\mathcal{T}'$  has no Aronszajn subtree, every point of  $\mathcal{T}'$  lies on an  $\omega_1$ -branch of  $\mathcal{T}'$ . It is now an easy matter to inductively (on the levels) embed  $\mathcal{T}$  into  $\mathcal{T}'$ .  $\square$

Let us return now to the property  $\Phi$ , and show how the failure of  $\Phi$  is closely connected with the existence of Aronszajn subtrees of trees. Let  $\Delta$  denote the following proposition, often referred to as Chang's Conjecture: If  $\mathcal{M} = \langle \omega_2, \omega_1, \dots \rangle$  is an arbitrary first-order structure with a countable language, there is  $\mathcal{N} = \langle B, B \cap \omega_1, \dots \rangle \prec \mathcal{M}$  such that  $|B| = \omega_1$  and  $|B \cap \omega_1| \leq \omega$ . Silver [3] has shown that  $\text{Con}(\text{ZFC} + \text{"there is a Ramsey cardinal"}) \rightarrow \text{Can}(\text{ZFC} + \text{GCH} + \Delta)$ . (And since  $\Delta$  implies the existence of Solovay's  $\aleph_1^\#$ , the large cardinal assumption here probably cannot be weakened very much, if at all.)

#### Theorem 7.

Assume  $\Delta + \text{GCH}$ . If  $\Phi$  fails, then there is an  $\omega_2$ -Aronszajn tree with no Aronszajn subtree.

Proof: Let  $\rho$  be an order type of cardinality  $\omega_2$  such that

$\omega_2, \omega_2^* \not\leq \rho$ , but for any  $\sigma \subseteq \rho$  of cardinality  $\omega_1$ , either  $\omega_1 \leq \sigma$  or else  $\omega_1^* \leq \sigma$ . Assume  $\rho = \langle \omega_2, - \rangle$  for definiteness.

Define, by induction on the levels, a tree,  $\mathcal{T}$ , as follows:  
The elements of  $\mathcal{T}$  are non-empty intervals of  $\rho$  and the ordering,  $<_{\mathcal{T}}$ , is  $\supset$ . Set  $T_0 = \{\rho\}$ . If  $I \in T_\alpha$  and  $|I| = 1$ ,  $I$  has no successors in  $\mathcal{T}$ . If  $I \in T_\alpha$  and  $|I| > 1$ , let  $\alpha_I$  be the least ordinal in  $I$  not maximal in  $I$  and let  $\{\xi \in I \mid \xi \rightarrow \alpha_I\}$  and  $\{\xi \in I \mid \alpha_I \rightarrow \xi\}$  be the successors of  $I$  in  $T_{\alpha+1}$ . Finally, if  $\lim(\delta)$  and  $\mathcal{T} \restriction \delta$  is defined, let  $T_\delta = \{nb \mid b \text{ is a } \delta\text{-branch of } \mathcal{T} \restriction \delta \text{ and } nb \neq \emptyset\}$ .

We linearly order each level  $T_\alpha$  by setting, for each  $I, J \in T_\alpha$ ,  $I <_\alpha J$  iff  $(\forall \xi \in I)(\forall \zeta \in J)(\xi \rightarrow \zeta)$ .

Suppose that for some  $\alpha < \omega_2$ ,  $|T_\alpha| = \omega_2$ .

Let  $\alpha$  be the least such. Then, clearly,  $\lim(\alpha)$ .

Let  $f: \mathcal{T} \restriction \alpha + 1 \leftrightarrow T_\alpha$ , and consider the structure

$\mathcal{A} = \langle \mathcal{T} \restriction \alpha + 1, T \restriction \alpha, T_\alpha, f, <_\alpha, <^* \rangle$ , where  $<^*$  is the lexicographic order on  $\mathcal{T} \restriction \alpha + 1$  induced by the  $<_\beta$ ,  $\beta < \alpha$ .

Now,  $|\mathcal{T} \restriction \alpha + 1| = \omega_2$  and  $|T \restriction \alpha| = \omega_1$  (by choice of  $\alpha$ ), so let

$\mathcal{L} = \langle X, X \cap T \restriction \alpha, \dots \rangle \prec \mathcal{A}$  with  $|X| = \omega_1$  and  $|X \cap T \restriction \alpha| = \omega$ ,

by assumption. Since

(i)  $\mathcal{A} \models (\forall x, y \in T_\alpha)(x <_\alpha y \rightarrow (\exists z \in T \restriction \alpha)(x \leq^* z \leq^* y))$

it follows that

(ii)  $\mathcal{L} \models (\forall x, y \in T_\alpha \cap X)(x <_\alpha y \rightarrow (\exists z \in T \restriction \alpha \cap X)(x \leq^* z \leq^* y))$

For each  $I \in T_\alpha \cap X$ , let  $\theta(I) \in I$  be arbitrary.

Then, for  $I, J \in T_\alpha \cap X$ ,  $I <_\alpha J$  iff  $\theta(I) \rightarrow \theta(J)$ .

Set  $\sigma = \{\theta(I) \mid I \in T_\alpha \cap X\}$ . Thus  $\sigma \subset \rho$ . And since

$f: X \leftrightarrow T_\alpha \cap X$ ,  $|\sigma| = \omega_1$ . We show that  $\omega_1, \omega_1^* \not\leq \sigma$ , a contradiction.

Suppose  $\omega_1 \leq \sigma$ . (The case  $\omega_1^* \leq \sigma$  is similar). It follows that

$\omega_1 \leq \langle T_\alpha \cap X, <_\alpha \restriction X^2 \rangle$ , so let  $\langle I_\nu \mid \nu < \omega_1 \rangle$  be a  $<_\alpha$ -increasing

sequence from  $T_\alpha \cap X$ . By (ii), we can inductively pick members  $J_\nu$

from  $T \restriction \alpha \cap X$ , for  $\nu = 0, \nu < \omega_1$ , so that  $\lim(\nu_1) \& \lim(\nu_2) \&$

$\nu_1 < \nu_2 \rightarrow I_{\nu_1} <^* J_{\nu_1} <^* I_{\nu_2}$ . But  $|T \restriction \alpha \cap X| = \omega$ , so we have a

contradiction.

Hence, for all  $\alpha < \omega_2$ ,  $|T_\alpha| \leq \omega_1$ .

Suppose  $T$  has an  $\omega_2$ -branch  $b = \langle I_\alpha \mid \alpha < \omega_2 \rangle$ . For each  $\alpha < \omega_2$  there is  $J_{\alpha+1} \in T_{\alpha+1}$  such that  $I_\alpha <_T J_{\alpha+1}$  and  $J_{\alpha+1} \neq I_{\alpha+1}$ . Let  $\theta_{\alpha+1} \in J_{\alpha+1}$  for each  $\alpha$ . Either  $|\{\theta_{\alpha+1} \mid \alpha < \omega_2 \text{ \& } I_{\alpha+1} <_{\alpha+1} J_{\alpha+1}\}| = \omega_2$  or else  $|\{\theta_{\alpha+1} \mid \alpha < \omega_2 \text{ \& } J_{\alpha+1} <_{\alpha+1} I_{\alpha+1}\}| = \omega_2$ . In the first case, the requisite  $\theta_{\alpha+1}$ 's form a  $\rightarrow$ -decreasing chain of type  $\omega_2$ , in the second case it is  $\rightarrow$ -increasing. So, in either event, we have a contradiction, since  $\omega_2, \omega_2^* \not\leq \rho$ . Hence  $T$  is an  $\omega_2$ -Aronszajn tree.

Clearly, any Aronszajn subtree of  $T$  will likewise correspond to a subtype  $\sigma \subseteq \rho$ ,  $|\sigma| = \omega_1$ , such that  $\omega_1, \omega_1^* \not\leq \sigma$ . Thus  $T$  cannot have an Aronszajn subtree, and we are done.  $\square$

Remark. By a simple generalisation of the proof of Theorem 6, one can construct, in ZFC, a normal  $(\omega_2, \omega_2)$ -tree with no Aronszajn (and no  $\omega_2$ -Aronszajn) subtree. We do not know if it is possible to construct, in ZFC + GCH, an  $\omega_2$ -Aronszajn tree with no Aronszajn subtree. If such were possible, however, we would immediately have a proof of  $\neg \Phi$  in ZFC + GCH, since the lexicographic ordering of such a tree is easily seen to provide a counterexample to  $\Phi$ . In view of an last result, this would seem to be the only hope of establishing  $\neg \Phi$  in ZFC + GCH. However, by  $\Delta$  itself, the more obvious sorts of  $\omega_2$ -Aronszajn trees which one can construct in ZFC + GCH do have Aronszajn subtrees, so this approach does not appear to be very hopeful. Much more likely, in our opinion, is that in Silver's model of  $\Delta$  (or perhaps a slight modification of it), every  $\omega_2$ -Aronszajn tree does have an Aronszajn subtree, whence, by theorem 7, we have at once the consistency of ZFC + GCH +  $\Phi$ . Unfortunately, a proof of this has so far eluded us.

Postscript

Since writing this paper, we obtained a proof of the result  
 $\text{Con}(\text{ZFC} + \text{"there is a Ramsey cardinal"}) \rightarrow \text{Con}(\text{ZFC} + 2^{\omega} = 2^{\omega_1} = \omega_2 + \Phi)$  .  
The proof will appear elsewhere. We still do not know if  $\Phi$  is  
consistent with GCH .

### References

1. K.J. Devlin. Aspects of Constructibility. Springer-Verlag:  
Lecture Notes in Mathematics [1973]
2. P. Erdős and A. Hajnal. Unsolved and Solved Problems in Set  
Theory. To appear.
3. J. Silver. The Consistency of Chang's Conjecture (unpublished).
4. J. Silver. The Independence of Kurepa's Conjecture.....  
AMS Proc. Symp. in Pure Maths. XIII, Part 1,  
pp. 383 - 390.